arXiv:0909.3771v1 [math.AG] 21 Sep 2009

WONDERFUL VARIETIES OF TYPE B AND C

P. BRAVI AND G. PEZZINI

Abstract. We show that the proof of Luna's conjecture about the classification of general wonderful G-varieties can be reduced to the analysis of finitely many families of primitive cases. We work out all primitive cases arising with any classical group G.

Luna's conjecture states that, for any semisimple complex algebraic group G, *wonderful G-varieties* are classified by *spherical systems* of G ([12]).

This has been proved for any G adjoint with simply laced Dynkin diagram ([12, 5, 1]) or under other special hypotheses ([13, 3]): one reduces to the case of *primitive* spherical systems, which can be listed, and shows that any primitive spherical system can be realized as the spherical system of a (unique) wonderful variety.

Afterwards, the uniqueness part of the conjecture has been proved in general, with different methods ([10]).

Here we follow the approach of [12]. The argument of reduction to the primitive spherical systems is adapted to the general adjoint (not necessarily simply laced) case. In other words, we show that, in order to complete the proof of the conjecture, it is enough to prove that there exists a wonderful variety with any given primitive spherical system.

The list of primitive spherical systems is given in [2].

Although it would be possible, we do not prove the existence of all the corresponding wonderful varieties. The list is very long and other general proofs of the conjecture seem now possible ([7, 8]).

We rather make use of further ideas coming from [10] and [4] to explain that it is possible to reduce the analysis to a smaller class of spherical systems: without *quotients of higher defect*. We also give a further special reduction argument to spherical systems without *tails*.

Finally, we analyze some relevant primitive spherical systems, mainly of type B and C. Together with [12, 5, 3] this completes the explicit case-by-case checking of Luna's conjecture for any G of classical type.

1. Wonderful varieties and spherical systems

1.1. **Basic definitions.** Let G be a semisimple complex algebraic group, T a maximal torus, B a Borel subgroup containing T, S the corresponding set of simple roots of the root system of (G,T). For all $S' \subset S$, $P_{S'}$ denotes the corresponding parabolic subgroup containing B (namely, S' is a set of simple roots of the root system of (L,T), where $L = L_{S'}$ is the Levi subgroup of $P_{S'}$ containing T). Similarly, $P_{-S'}$ denotes the opposite parabolic subgroup of $P_{S'}$, with respect to the maximal torus T.

Definition 1.1.1. A G-variety is called *wonderful* of rank r if it is smooth, complete, with r smooth prime G-divisors $D_1, \ldots, D_r$ with normal crossings, such that the G-orbit closures are exactly all the intersections $\cap_{i\in I} D_i$, for any $I \subseteq \{1, \ldots, r\}$.

We will always assume that the center of G acts trivially on a wonderful G-variety. This is justified by the results of Luna in [12], where the classification is reduced to the case where G is adjoint.

A wonderful G-variety is known to be projective and spherical, see [11].

Let X be a wonderful G-variety. There exists a unique point $z \in X$ stabilized by B_- (where B_- is the Borel subgroup opposite to B with respect to T, i.e. $B_- \cap B = T$). The orbit $G.z$ is the (unique) closed orbit, $\cap_{i=1}^r D_i$; the parabolic subgroup opposite to G_z will be denoted by P_X. Set S^p_X such that $P_X = P_{S^p_X}$.

The T-weights occurring in the normal space $T_zX/T_zG.z$ of $G.z$ at z are called spherical roots, their set is denoted by Σ_X. Spherical roots are in bijective correspondence with prime G-divisors, say $\sigma \mapsto D^\sigma$, and form a basis of the lattice of B-weights in $\mathbb{C}(X)$ (we are identifying B-weights with T-weights).

The not G-stable prime B-divisors of X are called colors, their set is denoted by Δ_X. The colors are representatives of a basis of Pic$X$, so one can define a $\mathbb{Z}$-bilinear pairing, called Cartan pairing, $c_X \colon \mathbb{Z}\Delta_X \times \mathbb{Z}\Sigma_X \to \mathbb{Z}$ such that $[D^\sigma] = \sum_{D\in\Delta_X} c_X(D,\sigma)[D]$.

For all $\alpha \in S$, set $\Delta_X(\alpha) = \{D \in \Delta_X : P_{\{\alpha\}}.D \neq D\}$. One has $\cup_{\alpha\in S}\Delta_X(\alpha) = \Delta_X$ and, for all $\alpha \in S$, card$(\Delta_X(\alpha)) \leq 2$. Clearly, $\alpha \in S^p_X$ if and only if $\Delta_X(\alpha) = \emptyset$. Moreover, if $D \in \Delta_X(\alpha)$ and card$(\Delta_X(\alpha)) = 1$ then $c_X(D,-)$ is uniquely determined by α. One has card$(\Delta_X(\alpha)) = 2$ if and only if $\alpha \in S \cap \Sigma_X$, in this case, say $\Delta_X(\alpha) = \{D^+, D^-\}$, $c_X(D^+,-)$ and $c_X(D^-,-)$ are not always determined by α, but their sum is. Let $\mathbf{A}_X$ denote the subset of colors $D \in \cup_{\alpha\in S\cap\Sigma_X}\Delta_X(\alpha)$ endowed with the $\mathbb{Z}$-linear functionals $c_X(D,-)$.

Definition 1.1.2. The datum of $(S^p_X, \Sigma_X, \mathbf{A}_X)$, also denoted by $\mathscr{S}_X$, is called the spherical system of X.

Any G-orbit closure X' of X is a wonderful G-variety itself. Its set of spherical roots $\Sigma' = \Sigma_{X'}$ is a subset of Σ_X, and X' is called the localization of X with respect to Σ' and denoted by $X' = X_{\Sigma'}$. The spherical system of X' is given by $S^p_{X'} = S^p_X$, $\Sigma_{X'} = \Sigma'$ and $\mathbf{A}_{X'}$, which can be identified with the subset of colors $D \in \cup_{\alpha\in S\cap\Sigma'}\Delta_X(\alpha)$; one has $c_{X'}(D,\sigma) = c_X(D,\sigma)$ for all $\sigma \in \Sigma'$.

In particular, any spherical root of X is the spherical root of a wonderful G-variety of rank 1. The wonderful G-varieties of rank 1 are well known, for all G. The finite set of spherical roots of wonderful G-varieties of rank 1 is denoted by $\Sigma(G)$.

The spherical system of X is determined by the spherical systems of all the localizations $X_{\Sigma'}$ of rank 2 (actually, it is enough to restrict to the localizations of rank 1 and those of rank 2 with a simple spherical root). Furthermore, the wonderful G-varieties of rank 2 are known, for all G.

1.2. **Spherical systems.**

A spherical system of G of rank $r \leq 2$ is, by definition, the spherical system of a wonderful G-variety of rank r. Wonderful G-varieties of rank $r \leq 2$ are classified by spherical systems of G of rank r.

Definition 1.2.1. A spherical system of G (of rank equal to card$\,\Sigma > 2$) is a triple $S^p \subset S$, $\Sigma \subset \Sigma(G)$ and $\mathbf{A} = \cup_{\alpha\in S\cap\Sigma}\mathbf{A}(\alpha)$, endowed with a $\mathbb{Z}$-bilinear pairing

$c\colon \mathbb{Z}\mathbf{A}\times\mathbb{Z}\Sigma\to\mathbb{Z}$, such that for all $\Sigma'\subset\Sigma$ with $\operatorname{card}\Sigma'=2$ the triple S^p, Σ' and $\mathbf{A}_{\Sigma'}=\cup_{\alpha\in S\cap\Sigma'}\mathbf{A}(\alpha)$, with the restriction of c, is the spherical system of a wonderful G-variety of rank 2.

Conjecture 1.2.1 ([12])**.** *Wonderful G-varieties are classified by spherical systems of G.*

2. Geometric constructions

2.1. **The set of all colors.** Given an abstract spherical system $\mathscr{S}=(S^p,\Sigma,\mathbf{A})$ for a given adjoint group G, it is possible to define a set $\Delta\supseteq\mathbf{A}$ of its colors (and a corresponding extension of c), which plays the role of Δ_X. Following [12] and its notations, we identify each element in $\Delta\setminus\mathbf{A}$ with the simple roots it is moved by. This gives a disjoint union:

$$\Delta=\mathbf{A}\cup\Delta^{a'}\cup\Delta^{b}$$

The set $\Delta^{a'}$ is the set of simple roots α such that 2α is a spherical root. For such a color D, we have $c(D,-)=\frac{1}{2}\langle\alpha^\vee,-\rangle$. The set Δ^b is:

$$\left(S\setminus\left(\Sigma\cup\frac{1}{2}\Sigma\cup S^p\right)\right)\Big/\sim$$

where $\alpha\sim\beta$ if $\alpha=\beta$, or if $\alpha\perp\beta$ and $\alpha+\beta\in\Sigma$. For such a color D, we have $c(D,-)=\langle\alpha^\vee,-\rangle$, for α any representative of the $\sim$-equivalence class associated to D.

2.2. **Localization and morphisms.** Let X be a wonderful variety and S' a subset of S. Consider a Levi subgroup $L_{S'}$ of the parabolic subgroup $P_{S'}$; the variety X contains a well defined wonderful $L_{S'}$-subvariety $X_{S'}$, called the *localization* of X in S' (see [12] for details). Its spherical system satisfies: $S^p_{X_{S'}}=S^p_X\cap S'$; the set $\Sigma_{X_{S'}}$ is the set of spherical roots of X whose support is contained in S'; the set $\mathbf{A}_{X_{S'}}$ is the union of $\Delta_X(\alpha)$ for all $\alpha\in S'\cap\Sigma_X$. The localization of an abstract spherical system $\mathscr{S}$ in S' is defined analogously.

Certain classes of morphisms between wonderful varieties can be represented by subsets of the colors of the domain. Let $f\colon X\to Y$ be a surjective G-equivariant morphism between two wonderful G-varieties, and define $\Delta_f=\{D\in\Delta_X\mid f(D)=Y\}$. Let now Δ' be a subset of Δ_X; we say that Δ' is *distinguished* if there exists a linear combination δ of elements of Δ' with positive coefficients, such that $c_X(\delta,\sigma)\geq 0$ for all spherical roots σ. We define the *quotient* spherical system $\mathscr{S}/\Delta'$ in the following way:

1. $S^p_X/\Delta'=\{\alpha\in S\mid\Delta(\alpha)\subseteq\Delta'\}$;
2. $\Sigma_X/\Delta'=$ the indecomposable elements (or equivalently the minimal generators) of the semigroup $\{\sigma\in\mathbb{N}\Sigma_X\mid c_X(\delta,\sigma)=0\ \forall\delta\in\Delta'\}$;
3. $\mathbf{A}_X/\Delta'=\bigcup_{\alpha\in S\cap\Sigma_X/\Delta'}\mathbf{A}(\alpha)$.

The bilinear pairing for $\mathscr{S}/\Delta'$ is induced from c_X in a natural way.

We say that Δ' is *(*)-distinguished*, if it is distinguished and the set Σ_X/Δ' is a basis of the $\mathbb{Z}$-module $\{\sigma\in\mathbb{Z}\Sigma_X\mid c_X(\sigma,\delta)=0\ \forall\delta\in\Delta'\}$.

Proposition 2.2.1 ([12])**.** *Let X be a wonderful G-variety. The map $f\mapsto\Delta_f$ induces a bijection between the set of (*)-distinguished subsets of Δ_X and the set of equivalence classes of couples (f,Y) where $f\colon X\to Y$ is a G-equivariant, surjective* *map with connected fibers, and where the equivalence relation is induced naturally by G-equivariant isomorphisms $Y \to Y'$.*

In principle, a distinguished but not (*)-distinguished set of colors corresponds to a G-equivariant morphism $X \to Y$ too. In this case Y satisfies the definition of a wonderful variety, except for the smoothness condition on Y and its prime G-divisors. Such a variety is also called the *canonical embedding* of its open G-orbit.

On the other hand, there is no known example of a distinguished set of colors that is not (*)-distinguished, and we conjecture indeed that none exists in general.

A (*)-distinguished subset $\Delta' \subseteq \Delta_X$ is

- *smooth* if $\Sigma_X/\Delta' \subseteq \Sigma_X$;
- *homogeneous* (or *parabolic*) if $\Sigma_X/\Delta' = \emptyset$.

Proposition 2.2.2 ([12])**.** *A G-equivariant, surjective map with connected fibers between wonderful varieties $f : X \to Y$ is smooth if and only if Δ_f is smooth. Moreover, Y is homogeneous if and only if Δ_f is homogeneous.*

All the above definitions can be obviously given also for an abstract spherical system $\mathscr{S}$.

Definition 2.2.1. Let $\mathscr{S}$ be a spherical system with set of colors Δ, and Δ_1, Δ_2 two subsets of Δ. We say that Δ_1 and Δ_2 *decompose* $\mathscr{S}$ if:

1. Δ_1 and Δ_2 are non-empty and disjoint;
2. Δ_1, Δ_2 and $\Delta_3 = \Delta_1 \cup \Delta_2$ are (*)-distinguished;
3. $(\Sigma \setminus (\Sigma/\Delta_1)) \cap (\Sigma \setminus (\Sigma/\Delta_2)) = \emptyset$;
4. $(S^p/\Delta_1) \setminus S^p$ and $(S^p/\Delta_2) \setminus S^p$ are orthogonal;
5. Δ_1 or Δ_2 are smooth.

Definition 2.2.2. Let $\mathscr{S}$ be a spherical system. A color $\delta \in \mathbf{A}$ is *projective* if $c(\delta, \sigma) \geq 0$ for all $\sigma \in \Sigma$.

Definition 2.2.3. A spherical system $\mathscr{S}$ is *primitive* if $\mathrm{supp}(\Sigma) = S$, there is no projective color, and there is no pair $\Delta_1, \Delta_2 \subset \Delta$ decomposing $\mathscr{S}$.

2.3. **Reduction to primitive spherical systems.**

To prove Conjecture 1.2.1 it is enough to show that there exists a (unique) wonderful G-variety with any given primitive spherical system. This is clear from the following:

Proposition 2.3.1. *Let $\mathscr{S} = (S^p, \Sigma, \mathbf{A})$ be a spherical system of G. If one of the following conditions is fullfilled then there exists a (unique) wonderful G-variety with $\mathscr{S}$ as spherical system.*

1. *There exist a proper subset S' of simple roots containing* $\mathrm{supp}(\Sigma)$ *and a (unique) wonderful $L_{S'}$-variety Y with spherical system equal to the localization of $\mathscr{S}$ with respect to S'.*
2. *There exist two (*)-distinguished subsets of colors Δ_1 and Δ_2 decomposing $\mathscr{S}$ and for $i = 1, 2, 3$ there exists a (unique) wonderful G-variety X_i with spherical system equal to the quotient of $\mathscr{S}$ by Δ_i, where $\Delta_3 = \Delta_1 \cup \Delta_2$.*
3. *There exist a projective element $\delta \in \mathbf{A}$ and a (unique) wonderful G-variety X_δ with spherical system equal to the quotient of $\mathscr{S}$ by $\{\delta\}$.*

In the first two cases Proposition 2.3.1 has been proved in [12, Section 3]. In the first case X is G-isomorphic to the parabolic induction $G *_{P_{S'}} Y$. In the second case X is G-isomorphic to the fiber product $X_1 \times_{X_3} X_2$.

In *loc.cit.* the third case has also been proved, but with some additional hypothesis: the support of δ is requested to intersect only the support of simple spherical roots. The latter is always true if G has a simply laced Dynkin diagram.

In §3.2 we prove it in full generality.

In *loc.cit.* it is also shown in general that, if X is a wonderful G-variety with $(S^p, \Sigma, \mathbf{A})$ as spherical system with a projective element $\delta \in \mathbf{A}$, then the morphism associated to δ is a projective fibration, namely smooth with fibers isomorphic to a projective space.

3. Wonderful subgroups

3.1. **Finding the wonderful subgroup.**

3.1.1. *Minimal inclusions.* Let $\mathscr{S}$ be a spherical system. In view of the following proposition, we say that $\mathscr{S}$ is *reductive* if there exists a linear combination σ of spherical roots with non-negative coefficients, such that $c(\delta, \sigma) > 0$ for all colors δ.

Proposition 3.1.1 ([12]). *Let X be a wonderful variety, with generic stabilizer H. Then H is reductive if and only if $\mathscr{S}$ is reductive.*

Let $\mathscr{S} = (\Sigma, S^p, \mathbf{A})$ be a spherical system with set of colors Δ, and suppose $\mathscr{S}$ is not reductive. This should correspond to a wonderful subgroup H, where moreover H is contained in some proper parabolic subgroup. Here and until the end of §3.1 we suppose that H exists.

Up to conjugating H if necessary, it follows that there exists a proper parabolic subgroup $Q_- \supseteq B_-$ containing H and minimal with respect to this property. It is possible to choose Levi subgroups L of H and L_{Q_-} of Q_- so that $L \subseteq L_{Q_-}$. From minimality of Q_- it follows that L is very reductive in L_{Q_-} and $H^r \subseteq Q_-^r$: in particular, $H^u \subseteq Q_-^u$ and the connected center C of L is contained in the connected center C_{Q_-} of L_{Q_-}.

There exists also a wonderful subgroup K such that $Q_- \supseteq K \supseteq H$ and K/H is connected, that is a wonderful subgroup properly containing H and minimal in the following sense (see [4], Section 2.3.5).

Choose a Levi component L_K of K in such a way that $L_{Q_-} \supseteq L_K \supseteq L$. Then the following holds:

1. the Levi subgroups L_K, L differ at most only for their connected centers;
2. we have $K^u \supset H^u \supseteq (K^u, K^u)$;
3. the quotient $\operatorname{Lie} K^u / \operatorname{Lie} H^u$ is simple as L-module.

These subgroups Q_- and K correspond resp. to two quotients $\mathscr{S}/\Delta_{Q_-}$ and $\mathscr{S}/\Delta_K$ of $\mathscr{S}$.

Let us now abandon the hypothesis that H exists. We usually find Δ_{Q_-} and Δ_K which play the same role:

- Δ_{Q_-}, a minimal homogeneous subset of colors;
- Δ_K, a special (not any) minimal (*)-distinguished subset of colors contained in Δ_{Q_-}, such that the wonderful subgroup K corresponding to $\mathscr{S}/\Delta_K$ is known.

We then find H inside K, as above. Assuming $(L, L) = (L_K, L_K)$, it remains to find C and H^u.

3.1.2. *The connected center.* The dimension of C is immediately given by the well known formula:

$$\dim C = \operatorname{card}\Delta - \operatorname{card}\Sigma.$$

This integer is also called $\mathrm{d}(\mathscr{S})$, *defect* of $\mathscr{S}$.

Consider all B-proper rational functions on G/H having neither zeros nor poles along colors in Δ_{Q_-}. Their B-weights form a sublattice $\mathscr{N}$ of $\mathbb{Z}\Sigma$: fix a basis $\mathscr{B}$ of it.

For each color D in $\Delta\setminus\Delta_{Q_-}$ define α_D as the unique simple root moving D (this root is unique because Δ_{Q_-} is homogeneous). Define also λ_D to be the fundamental dominant weight associated to α_D, and $\lambda^*_D = -w_0(\lambda_D)$, where w_0 is the longest element in the Weyl group.

Finally, for all $\gamma\in\mathscr{N}$, define $\lambda(\gamma)$ to be the weight:

$$\sum_{D\in\Delta\setminus\Delta_{Q_-}} c(D,\gamma)\cdot\lambda^*_D$$

restricted to the subgroup C_{Q_-}.

Lemma 3.1.1. *The subgroup C is the connected part of the intersection of the kernels of the weights $\lambda(\gamma)$, for γ ranging through $\mathscr{N}$ (or, equivalently, $\mathscr{B}$).*

Proof. Let $\gamma\in\mathscr{N}$ and consider a B-proper rational function f_γ on G/H having B-weight γ. Call F_γ the pull-back of f_γ on G along the projection $G\to G/H$.

Since f_γ has neither zeros nor poles on colors of Δ_{Q_-}, F_γ has neither zeros nor poles on the pull-backs on G of these colors. Its zeros or poles must then lie only on the pullbacks on G of colors of G/Q_- along the projection $G\to G/Q_-$.

This means that F_γ is Q_--proper under right translation, and it is immediate to see that the Q_--weight of F_γ is just $\lambda(\gamma)$. But F_γ is also of course H-stable under right translation, and this implies that $\lambda(\gamma)$ is constantly 1 on C.

Vice versa, suppose to have any weight χ of C_{Q_-} which is constantly 1 on C: considering χ as a Q_--weight, we can write it as a linear combination with integer coefficients of weights of the form λ^*_D for D ranging through the colors of G/Q_-.

So χ is the Q_--weight (under right translation) of some rational function on G, call it F, which is B-proper under left translation and Q_--proper under right translation, whose zeros and poles lie on the pullbacks on G of colors of G/Q_- along the projection $G\to G/Q_-$.

The function F under right translation is Q^u_--stable and (L_{Q_-},L_{Q_-})-stable; it is C_{Q_-}-proper but C-stable. Thus F is H-stable, and it descends to a rational function f on G/H. Having neither zeros nor poles along colors in Δ_{Q_-} by construction, the function f is equal to f_γ for some $\gamma\in\mathscr{N}$. □

3.1.3. *The unipotent radical.* The unipotent radical H^u is a subgroup of K^u, containing (K^u,K^u) and such that $\operatorname{Lie}K^u/\operatorname{Lie}H^u$ is a simple L-module. For computations in particular cases, it is useful to recall that H^u is also a subgroup of Q^u_-, and that the quotient $\operatorname{Lie}Q^u_-/\operatorname{Lie}H^u$ is a spherical L-module. The following lemma follows from the results of [10].

Lemma 3.1.2 ([10])**.** *The spherical system $\mathscr{S}$ uniquely determines the structure of $\operatorname{Lie}Q^u_-/\operatorname{Lie}H^u$ as an L-module.*

Let us describe how to calculate the L-module structure of $\operatorname{Lie}Q^u_-/\operatorname{Lie}H^u$ starting from the spherical system $\mathscr{S}$. A fiber $\mathscr{F}$ of the morphism $G/H\to Q_-$ is affine and isomorphic to $L_{Q_-}\times_L V$ where $V=\mathrm{Lie}\,Q_-^u/\mathrm{Lie}\,H^u$. Restricting B-proper rational functions on G/H to $\mathscr{F}$ we obtain $(B\cap L_{Q_-})$-proper rational functions: this induces a bijection between the associated lattices of B- (resp. $(B\cap L_{Q_-})$-) weights (a proof of this fact can be found in [6]).

Restricting $(B\cap L_{Q_-})$-proper regular functions from $\mathscr{F}$ to V, we obtain all $\mathbb{C}[V]^{(B\cap L)}$. On the other hand, regular functions on $\mathscr{F}$ come exactly from rational functions on G/H having no poles on colors in Δ_{Q_-}.

Therefore the weight monoid of V as a spherical L-module is given by restricting to $B\cap L$ all B-weights in Ξ that are non-negative on Δ_{Q_-}. Finally, the classification of spherical modules is used to determine the L-module structure of V.

3.2. **Projective fibrations.** We proceed completing the proof of Proposition 2.3.1. Let $\mathscr{S}=(S^p,\Sigma,\mathbf{A})$ be a spherical system with a projective $\delta\in\mathbf{A}$. Set $\mathrm{supp}\,\delta=\{\alpha\in S:\delta\in\mathbf{A}(\alpha)\}$.

Lemma 3.2.1. *Suppose* $\mathrm{supp}\,\delta\cap\mathrm{supp}(\Sigma\setminus S)\neq\emptyset$. *If there exists a wonderful variety X_δ with spherical system $\mathscr{S}/\{\delta\}$, then there exists a wonderful variety with spherical system $\mathscr{S}$.*

Recall that (as said in §2.3) the assertion is true also if $\mathrm{supp}\,\delta\cap\mathrm{supp}(\Sigma\setminus S)=\emptyset$.

Let us first prove that one can reduce to the case of $\mathrm{card}(\mathrm{supp}\,\delta)=1$, so we assume the assertion in this case.

Let X be a wonderful variety with spherical system $\mathscr{S}$ with a projective δ with $\mathrm{card}(\mathrm{supp}\,\delta)=1$. Let K be the generic stabilizer of X_δ, with Levi decomposition $K=L_K\,K^u$. Then a generic stabilizer $H\subset K$ of X has the same Levi factor $L=L_K$ and unipotent radical H^u, such that $\mathrm{Lie}\,H^u$ is a 1-codimensional L-submodule of $\mathrm{Lie}\,K^u$.

Let $\mathscr{S}$ be a spherical system with a general projective element δ. Let X_δ be a wonderful variety with spherical system $\mathscr{S}/\{\delta\}$, and let K be its generic stabilizer with Levi decomposition $K=L_KK^u$. Then for each $\alpha\in\mathrm{supp}\,\delta$ we can consider the spherical system obtained from $\mathscr{S}/\{\delta\}$ by adding the simple spherical root α such that one of the two corresponding elements of $\mathbf{A}(\alpha)$, say δ_α^+ is projective, hence $\mathrm{card}(\mathrm{supp}\,\delta_\alpha^+)=1$. Then there exists a wonderful variety with this spherical system and generic stabilizer given by $H_\alpha=L_KH_\alpha^u$ where $\mathrm{Lie}\,H_\alpha^u$ is a 1-codimensional L_K-submodule in $\mathrm{Lie}\,K^u$. Take the subgroup $H=L\,H^u$ given as follows: $\mathrm{Lie}\,H^u$ equal to a 1-codimensional (L_K,L_K)-submodule in $\mathrm{Lie}\,K^u$ and in general position among those containing

$$\bigcap_{\alpha\in\mathrm{supp}\,\delta}\mathrm{Lie}\,H_\alpha^u;$$

L equal to the normalizer of H^u in L_K. Then H is the generic stabilizer of a wonderful variety with spherical system $\mathscr{S}$.

Let now $\mathscr{S}$ be a spherical system with a projective element δ with $\mathrm{card}(\mathrm{supp}\,\delta)=1$, say $\mathrm{supp}\,\delta=\{\alpha\}$. If there exists a (*)-distinguished subset of colors Δ_2 disjoint from $\mathbf{A}(\alpha)$, then the subsets of colors $\Delta_1=\{\delta\}$ and Δ_2 decompose the spherical system $\mathscr{S}$. Therefore, by using the second case of Proposition 2.3.1, we can suppose that there exists no (*)-distinguished subset of colors Δ_2 disjoint from $\mathbf{A}(\alpha)$. This implies that the spherical system is indecomposable.

Clearly, we can also suppose that the spherical system is cuspidal.

Furthermore, we assumed $\mathrm{supp}\,\delta\cap(\Sigma\setminus S)\neq\emptyset$.

A careful combinatorial analysis then shows that the only possibilities are the following “minimal” spherical systems, which will be treated case by case in §4.4.

Here (and often in the rest of the paper) we describe spherical systems by providing their Luna diagrams, see [4] for their definition.

(1) G of type C_n, $n \geq 2$



(2) G of type G_2



(3) G of type F_4



(4) G of type F_4



(5) G of type F_4



(6) G of type B_n, $n \geq 3$



3.3. **Quotients of higher defect.** The above discussion about projective colors, supposing $\mathrm{card}(\mathrm{supp}\,\delta) > 1$, can be considered as a particular case of a more general situation. Let Δ_K be a minimal (*)-distinguished subset of colors of $\mathscr{S}$, and recall the notation of §3.1.1. We make the crucial assumption that $\mathrm{d}(\mathscr{S}/\Delta_K) > \mathrm{d}(\mathscr{S})$, namely $C < C_K$.

One can then determine the unipotent radical H^u by means of some localizations of $\mathscr{S}$. The idea is as follows. Suppose once again that H exists (with $C < C_K$). The unipotent radical K^u is of course stable under conjugation by C_K, and the same holds resp. for H^u and C. But H must be finite in its normalizer, hence no element of $C_K \setminus C$ is allowed to normalize H^u, otherwise it would be inside $\mathrm{N}_G H$.

Now we want to discuss $\mathrm{Lie}\, K^u/[\mathrm{Lie}\, K^u, \mathrm{Lie}\, K^u]$ together with its L-submodule $\mathrm{Lie}\, H^u/[\mathrm{Lie}\, K^u, \mathrm{Lie}\, K^u]$ under this point of view. The former must be a sum of L_K-modules:

$$\mathrm{Lie}\, K^u/[\mathrm{Lie}\, K^u, \mathrm{Lie}\, K^u] = V \oplus W$$

where W is a sum:

$$W = W_0 \oplus W_1 \oplus \ldots \oplus W_k$$

such that:

(3.3.1) the difference $\mathrm{d}(\mathscr{S}/\Delta_K) - \mathrm{d}(\mathscr{S})$ is equal to k;

(3.3.2) the modules $W_0, \ldots, W_k$ are all simple and isomorphic as L-modules; the action of L_K differs on them only for C_K;

(3.3.3) if C_K acts resp. with weights $\chi_0, \ldots, \chi_k$, the connected component of the subgroup where $\chi_0, \ldots, \chi_k$ coincide is exactly C;

(3.3.4) $\mathrm{Lie}\, H^u/[\mathrm{Lie}\, K^u, \mathrm{Lie}\, K^u]$ is a submodule of codimension $\dim W_i$ and in general position among those containing V.

In other words $\operatorname{Lie} H^u$ contains all $\operatorname{Lie} K^u$ except for a certain part W, of which $\operatorname{Lie} H^u$ contains only a "diagonal" submodule.

The submodules W_i can be found studying the geometry of the wondeful embedding X of G/H. Call $x \in X$ the point having H as its stabilizer, and choose a 1-parameter subgroup η of C_K. Any point on X of the form $\eta(t)x$, for $t \in \mathbb{C}^*$, has stabilizer $\eta(t)H\eta(t)^{-1}$. Consider now the limit:

$$x(\eta) = \lim_{t\to 0} \eta(t)x$$

and its stabilizer $H(\eta)$ in G. Depending on η, this point may lie in a non-open orbit; if we fix $i \in \{0, \ldots, k\}$ and suppose that η satisfies:

$$\langle \eta, \chi_i \rangle = 1, \text{ and } \langle \eta, \chi_j \rangle = 0 \ \forall j \neq i$$

then $x(\eta)$ lies evidently on a codimension 1 orbit of X, the group $H(\eta)$ has a Levi factor whose connected center $C(\eta)$ satisfies $C_K \supseteq C(\eta) \supset C$ and $\dim C(\eta) = \dim C + 1$, and finally:

$$\operatorname{Lie} H(\eta)^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u] \supset W_j \text{ iff } j = i.$$

This gives a way to find H^u starting from the stabilizers of points of X in codimension 1 orbits.

Let us then rephrase the above procedure without appel to the existence of H.

Let $\mathscr{S}$ be a spherical system with Δ_K as above. For any $\gamma \in \Sigma$ define $\Delta_K(\gamma)$ to be Δ_K restricted to the localization $\mathscr{S}_{\Sigma\setminus\{\gamma\}}$ of $\mathscr{S}$ with respect to $\Sigma \setminus \{\gamma\}$. Then there exist $k+1$ spherical roots $\gamma_0, \ldots, \gamma_k$ of $\mathscr{S}$ such that for all $i = 0, \ldots, k$:

(3.3.5) $\Delta_K(\gamma_i)$ is a minimal (*)-distinguished subset of colors of $\mathscr{S}_{\Sigma\setminus\{\gamma_i\}}$;

(3.3.6) we have:

$$\mathrm{d}\left(\frac{\mathscr{S}_{\Sigma\setminus\{\gamma_i\}}}{\Delta_K(\gamma_i)}\right) - \mathrm{d}(\mathscr{S}_{\Sigma\setminus\{\gamma_i\}}) = k - 1;$$

(3.3.7) $\operatorname{supp}(\Sigma \setminus \{\gamma_i\}) \neq S$.

Let us also suppose that the spherical systems $\mathscr{S}_{\Sigma\setminus\{\gamma_i\}}$ (for all i) and $\mathscr{S}/\Delta_K$ correspond to wonderful varieties having generic stabilizers resp. H_i and K (with $H_i \subset K$). Then:

(3.3.8) Levi subgroups L_i of H_i and L_K of K can be chosen in such a way that $(L_i, L_i) = (L_K, L_K)$ for all i;

(3.3.9) with this choice, the connected centers C_i of L_i are such that their connected intersection is equal to C as described in Lemma 3.1.1;

(3.3.10) each unipotent radical H_i^u contains (K^u, K^u), in such a way that the quotients $\operatorname{Lie} H_i^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u]$ are all isomorphic as (L_K, L_K)-modules;

(3.3.11) there exist L_K-submodules $V, W_1, \ldots, W_k$ of $\operatorname{Lie} K^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u]$, with all the W_i isomorphic as $C(L_K, L_K)$-modules, satisfying the conditions (3.3.2) and (3.3.3) above and such that $\operatorname{Lie} H_i^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u]$ (of codimension $\dim W_i$ in $\operatorname{Lie} K^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u]$) is in general position among those containing V and W_i, for all $i = 0, \ldots, k$.

Define C to be the connected intersection of the C_i's, and L to be $C(L_K, L_K)$. Define H^u in such a way that $\operatorname{Lie} H^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u]$ has codimension $\dim W_i$ in $\operatorname{Lie} K^u / [\operatorname{Lie} K^u, \operatorname{Lie} K^u]$ and is in general position containing V; and $H = L\,H^u$. Then:

(3.3.12) H is wonderful and its spherical system is $\mathscr{S}$.

Notice that once H is known to be wonderful, its spherical system is uniquely determined: let us use the subgroups $H(\eta)$ defined above, where η is a 1-parameter subgroup of C_K. As before, we can choose $\eta_0, \ldots, \eta_k$ such that $H(\eta_i) = H_i$. It follows that the spherical system of H has $k+1$ localizations of rank $r-1$ equal to the spherical systems of H_i.

We claim that all the previous statements hold for any spherical system $\mathscr{S}$ with a quotient of higher defect $\mathscr{S}/\Delta_K$. When $\mathscr{S}$ is primitive, they can be explicitly checked case by case, obtaining the following:

Proposition 3.3.1. *The existence for primitive spherical systems with quotients of higher defect is a consequence of the existence for primitive sperical systems without quotients of higher defect.*

3.4. **Tails.** Another general procedure applies to all spherical systems $\mathscr{S}$ obtained adding one of the following spherical roots γ to another spherical system $\mathscr{S}'$ (i.e. $\mathscr{S}'$ is the localization of $\mathscr{S}$ in $S' \subset S$, with $S' \cup \operatorname{supp}\gamma = S$). We always have (or ask) that γ glues on a unique connected component of S' of type A.

If $S' \cap \operatorname{supp}\gamma = \emptyset$ (then there exists a unique spherical root γ' non-orthogonal to $\operatorname{supp}\gamma$):

$b(m)$: if $m > 1$:



support of type B_m, $\gamma = \alpha_{p+1} + \ldots + \alpha_{p+m}$, with all roots α_{p+i} in S^p except for α_{p+1}; if $m = 1$:



support of type A_1, $\gamma = \alpha \in S \setminus S^p$ with α the short simple root of a component of S of type B, $c(D^+_\alpha, \gamma') = c(D^-_\alpha, \gamma') = -1$.

$b'(m)$: if $m > 1$:



support of type B_m, $\gamma = 2\alpha_{p+1} + \ldots + 2\alpha_{p+m}$, with all roots α_{p+i} in S^p except for α_{p+1}; if $m = 1$:

support of type A_1, $\gamma = 2\alpha$, $\alpha \in S \setminus S^p$ with α the short simple root of a component of S of type B.

$d(m)$: if $m > 2$:



support of type D_m, $\gamma = 2\alpha_{p+1} + \ldots + 2\alpha_{p+m-2} + \alpha_{p+m-1} + \alpha_{p+m}$, with all roots α_{p+i} in S^p except for α_{p+1}; if $m = 2$:



support of type $\mathsf{A}_1 \times \mathsf{A}_1$, $\gamma = \alpha + \alpha'$, $\alpha, \alpha' \in S \setminus S^p$ with α, α' at the "end" of a component of S of type D (i.e. $\{\beta \in S : \alpha \not\perp \beta\} = \{\beta \in S : \alpha' \not\perp \beta\}$).

If $S' \cap \operatorname{supp}\gamma \neq \emptyset$:

$c^*(m)$: for $m \geq 2$:



support of type C_m, $\gamma = \alpha_{p+1} + 2\alpha_{p+2} + \ldots + 2\alpha_{p+m-1} + \alpha_{p+m}$, with all roots α_{p+i} in S^p except for α_{p+1} and α_{p+2}, and $S' \cap \operatorname{supp}\gamma = \{\alpha_{p+1}\}$. In this case we will say that γ is *overlapping*.

Definition 3.4.1. A spherical system $\mathscr{S}$ has a *tail* in one of its spherical roots γ, if γ is one of the above cases $b(m), b'(m), d(m), c^*(m)$ and there exists a (*)-distinguished subset of colors Δ' such that the quotient $\mathscr{S}/\Delta'$ has rank 1 with γ as spherical root and $S^p = \{\beta \in S : \gamma \perp \beta\}$.

The set of colors Δ' with the above property always exists if $\operatorname{supp}\gamma$ is a connected component of S (so γ is necessarily of type $c^*(m)$): it is given by all the colors of $\mathscr{S}'$. Otherwise, if Δ' exists, it is given by all the colors of $\mathscr{S}'$ except one.

Being of rank 1, the wonderful variety associated to $\mathscr{S}/\Delta'$ exists: we call its generic stabilizer $M \subset G$. If our strategy is based on an induction on the number of tails, we can suppose the same for the wonderful variety associated to $\mathscr{S}'$: we call its generic stabilizer $K \subset L_{S'}$. We will now see how these two subgroups determine a candidate subgroup H for the spherical system $\mathscr{S}$.

3.4.1. *Tail of type $b(m)$.* An example of this kind of tail is the rank 2 spherical system of $\mathrm{SO}(2p+2m+1)$:



$$\mathscr{S} = (\{\alpha_2, \ldots, \alpha_{p-1}, \alpha_{p+2}, \ldots, \alpha_{p+m}\}, \{\alpha_1 + \ldots + \alpha_p, \alpha_{p+1} + \ldots + \alpha_{p+m}\}, \emptyset)$$

Its wonderful subgroup has Levi factor $\mathrm{GL}(p) \times \mathrm{SO}(2m)$ and Lie algebra of the unipotent radical equal to $\mathbb{C}^p \otimes \mathbb{C}^{2m} \oplus \wedge^2\mathbb{C}^p$.

The quotient $\mathscr{S}/\Delta'$ is the rank 1 case:

$$(\{\alpha_1, \ldots, \alpha_{p-1}, \alpha_{p+2}, \ldots, \alpha_{p+m}\}, \{\alpha_{p+1} + \ldots + \alpha_{p+m}\}, \emptyset)$$

whose wonderful subgroup is $(\mathrm{GL}(p) \times \mathrm{SO}(2m))Q_-^r$, and it is a parabolic induction by means of the parabolic subgroup $Q_- = P_{-S\setminus\{\alpha_p\}}$. The set S' is $\{\alpha_1, \ldots, \alpha_p\}$, and the localization $\mathscr{S}'$ is:

$$\mathscr{S}' = (\{\alpha_1 + \ldots + \alpha_p\}, \{\alpha_2, \ldots, \alpha_{p-1}\}, \emptyset)$$

whose wonderful subgroup is $\mathrm{GL}(p)$ inside $\mathrm{SL}(p+1)$. The morphism $\phi_{\Delta'}$ restricted to $\mathscr{S}'$ corresponds to the inclusion $\mathrm{GL}(p) \subset R_-$ where R_- is the maximal parabolic subgroup of $\mathrm{SL}(p+1)$ associated to $S' \setminus \{\alpha_p\}$.

Now let us discuss a general case. First of all $\operatorname{supp}\gamma$ is never a connected component of the Dynkin diagram of G: let β_0 be the unique $\beta_0 \in S \setminus \operatorname{supp}\gamma$ with $\beta_0 \not\perp \operatorname{supp}\gamma$. Recall also that the connected component of the Dynking diagram of $L_{S'}$ containing β_0 is of type A.

Let $m = \operatorname{card}(\operatorname{supp}\gamma)$. The system $\mathscr{S}/\Delta'$ is a parabolic induction of the wonderful $\mathrm{SO}(2m+1)$-variety having generic stabilizer $\mathrm{SO}(2m)$, where the induction is done using the parabolic subgroup Q_- associated to all simple roots except β_0. Therefore Q_- has a Levi part L_{Q_-} which contains the factor $\mathrm{SO}(2m+1)$: let us write $L_{Q_-} = \widetilde{L_{Q_-}} \times \mathrm{SO}(2m+1)$. With these notations of course $M = (\widetilde{L_{Q_-}} \times \mathrm{SO}(2m))Q_-^r$.

On the other hand, we can look at the quotient by Δ' *on* the localization $\mathscr{S}'$: it follows that $\mathscr{S}'$ itself has a quotient which is the rank 0 variety having generic stabilizer equal to the maximal parabolic subgroup $R_- \subset G^{S'}$ associated to $S' \setminus \beta_0$.

Thus K is contained in R_-, whose Levi part we denote by L_{R_-}. Choose a Levi part L_K of K contained in L_{R_-}.

If $\mathscr{S}'$ is not a parabolic induction (e. g. if $\mathscr{S}'$ is primitive), K does not contain R^u_-: in other words the L_K-module $\mathrm{Lie}\,R^u_-/(\mathrm{Lie}\,R^u_-\cap\mathrm{Lie}\,K^u)$ is non-trivial. It is also easy to see that $\mathrm{Lie}\,K^u=(\mathrm{Lie}\,R^u_-\cap\mathrm{Lie}\,K^u)+(\mathrm{Lie}\,L_{R_-}\cap\mathrm{Lie}\,K^u)$.

By construction L_{R_-} can be identified with $\widetilde{L_{Q_-}}$: we can set $L=L_K\times\mathrm{SO}(2m)$ as a subgroup of M. Moreover, the L_K-module $\mathrm{Lie}\,R^u_-\cap\mathrm{Lie}\,K^u$ corresponds evidently to an L-submodule $\mathrm{Lie}\,U$ of $\mathrm{Lie}\,Q^u_-$, in such a way that U contains (Q^u_-,Q^u_-), that $\mathrm{SO}(m)$ acts trivially on Q^u_-/U and the two quotients $\mathrm{Lie}\,R^u_-/(\mathrm{Lie}\,R^u_-\cap\mathrm{Lie}\,K^u)$, $\mathrm{Lie}\,Q^u_-/\mathrm{Lie}\,U$ are isomorphic as L_K-modules.

The candidate H^u is now $U(L_{R_-}\cap K^u)$, and we can set $H=L\,H^u$.

3.4.2. *Tail of type $b'(m)$.* This case is similar to the preceeding, the only difference is that we consider $\mathrm{N}_{\mathrm{SO}(2m+1)}\mathrm{SO}(2m)$ instead of $\mathrm{SO}(2m)$.

3.4.3. *Tail of type $c^*(m)$.* When $\mathrm{supp}\,\gamma$ is not a connected component of the Dynkin diagram of G, this case is similar to $b(m)$: the only difference is in the definition of L, where one must only use $\mathrm{Sp}(2m-2)$ inside $\mathrm{Sp}(2m)$, instead of $\mathrm{SO}(2m)$ inside $\mathrm{SO}(2m+1)$. More precisely, we have $L_{Q_-}=\widetilde{L_{Q_-}}\times\mathrm{Sp}(2m-2)\times\mathrm{SL}(2)$, and $L_{R_-}=\widetilde{L_{R_-}}\times\mathrm{SL}(2)$, where the second factor has simple root the first root of $\mathrm{supp}\,\gamma$. Here $\widetilde{L_{R_-}}$ can be identified by construction with $\widetilde{L_{Q_-}}$, and we set $L=L_K\times\mathrm{Sp}(2m-2)$.

When $\mathrm{supp}\,\gamma$ is a connected component of the Dynkin diagram of G, then $\mathscr{S}/\Delta'$ is not a parabolic induction: it is the wonderful $\mathrm{Sp}(2m)$-variety with generic stabilizer $\mathrm{Sp}(2m-2)\times\mathrm{SL}(2)$. The definition of H proceeds as before, with the only difference that there is no part U.

3.4.4. *Tail of type $d(m)$.* This case is similar to $b(m)$: we use $\mathrm{SO}(2m-1)$ inside $\mathrm{SO}(2m)$, instead of $\mathrm{SO}(2m)$ inside $\mathrm{SO}(2m+1)$.

3.4.5. *Proof of existence in tail cases.*

Proposition 3.4.1. *The existence for tail primitive spherical systems is a consequence of the existence for non-tail primitive sperical systems.*

Proof. We begin by excluding for a while tails of type $b(m)$. Let then $\mathscr{S}$ be a primitive spherical system with only tails of type $b'(m)$, $c^*(m)$ and $d(m)$.

We proceed by induction on the number of tails, so select a tail γ and let H be the candidate subgroup determined in the previous subsections. Let us also use all the elements introduced above which depend on the choice of γ: the localization $\mathscr{S}'$, the set of colors Δ', and the subgroups M, K, Q_-, R_-.

The combinatorial characterization of the group of G-equivariant automorphisms given in [10] says that if $\mathscr{S}$ is associated to a wonderful variety, then the latter has trivial such group. This because there is no tail of type $b(m)$. The same statement holds for $\mathscr{S}'$ too.

Therefore $\mathrm{N}_{G^{S'}}K=K$; from this and from the definition of H it follows that $\mathrm{N}_G H=H$. The subgroup H is hence wonderful.

We must now prove that the spherical system $\mathscr{S}_X=(S^p_X,\Sigma_X,\mathbf{A}_X)$ of the wonderful embedding X of G/H is equal to $\mathscr{S}$. First of all notice that L_K is a factor of L: call $\widetilde{L}$ the complementary factor of L; it is easy to see from the definition of H that it can be chosen in such a way that H is B-spherical and $\widetilde{L}$ is $(B\cap L_{\mathrm{supp}\,\gamma})$-spherical.

Let us call β_1, β_2 (and β_3) resp. the unique simple root of $\operatorname{supp}\gamma$ not orthogonal to $S \setminus \operatorname{supp}\gamma$ and the simple root(s) moving the color whose functional is positive on γ.

Our first claim is that $\gamma \in \Sigma_X$. To prove it, first consider the case $(\operatorname{supp}\gamma) \setminus \{\beta_1, \beta_2, \beta_3\} = \emptyset$. Here the tail is $b'(1), c^*(2)$ or $d(2)$ and the combinatorics of spherical systems implies easily that $\gamma \in \Sigma_X$, using the fact that $\mathrm{N}_G H = H$, that the variety X is not a parabolic induction, that the subgroup H is contained in M and that we have $\mathrm{d}(\mathscr{S}_X) = \mathrm{d}(\mathscr{S}')$ for these kinds of tail.

Now suppose $(\operatorname{supp}\gamma) \setminus \{\beta_1, \beta_2, \beta_3\} \neq \emptyset$. Call X' the wonderful embedding of G/M: its spherical system is a quotient of $\mathscr{S}_X$ thanks to the inclusion $H \subset M$.

Consider also the localization X'' of X' in $\operatorname{supp}\gamma$. It is the wonderful $L_{\operatorname{supp}\gamma}$-variety having spherical system $((\operatorname{supp}\gamma) \setminus \{\beta_2, \beta_3\}, \{\gamma\}, \emptyset)$: its generic stabilizer is (L_M, L_M). If we conjugate (L_M, L_M) with an element of $U_{-\alpha}$ for $\alpha \in (\operatorname{supp}\gamma) \setminus \{\beta_1, \beta_2, \beta_3\}$ then it remains a $(B \cap L_{\operatorname{supp}\gamma})$-spherical subgroup, where $U_{-\alpha}$ is the 1-dimensional unipotent subgroup of G associated to $-\alpha$.

Given the definition of H, this ensures that if we conjugate H with the same element of $U_{-\alpha}$ then it remains B-spherical: in other words $(\operatorname{supp}\gamma) \setminus \{\beta_1, \beta_2, \beta_3\} \subseteq S^p_X$. This implies immediately that $\gamma \in \Sigma_X$, thanks to a quick check on the lists of spherical roots in types $\mathsf{B}, \mathsf{C}, \mathsf{D}$ and the related allowed sets S^p (and again the fact that X is not a parabolic induction, $H \subset M$ and $\mathrm{N}_G H = H$).

Let us now consider the localization Y of X on S'. Our second claim is that this localization has spherical system $\mathscr{S}'$. To prove it, consider the G-equivariant map from X to the wonderful embedding of G/M: restricted to Y, it is a map onto $L_{S'}/R_-$. From the definition of H, it is obvious that the latter map and the map from the wonderful variety of $\mathscr{S}'$ to $L_{S'}/R_-$ have fibers with the same open R_--orbit: it follows that Y has spherical system $\mathscr{S}'$.

This finishes the proof if we exclude tails of type $b(m)$. Now let $\mathscr{S}$ be a primitive spherical system with k tails of type $b(m)$ (with possibly different values of m of course), and call again H the candidate subgroup given in the previous subsections. Call $\gamma_1, \ldots, \gamma_k$ the k tails of type $b(m)$, and define $\overline{\mathscr{S}}$ the same spherical system as $\mathscr{S}$ with each γ_i replaced by $2\gamma_i$.

By the first part of the proof, $\overline{\mathscr{S}}$ is the spherical system of a wonderful variety $\overline{X}$; by the definitions of the candidate generic stabilizers given in the previous subsections it is clear that the generic stabilizer of $\overline{X}$ is equal to $\mathrm{N}_G H$, and that $[\mathrm{N}_G H : H] = 2^k$. Now the subgroup $\mathrm{N}_G H$ is spherically closed, and we conclude thanks to the description given by Luna in [12] of spherical subgroups having a fixed spherical closure. □

4. Explicit computations

4.1. **Reductive cases.** We discuss the reductive primitive cases first. They are in some sense the "basic" cases in view of our approach via morphisms, see §3.1.

For G a product of classical groups of type A and D, we refer to the published work [5]. Table 1 contains the list of reductive wonderful subgroups of products of classical groups where at least one factor has type B or C (some of them have been already analyzed in [13] and [3]). A more explicit description of the spherical systems here represented by diagrams can be found in [2].

Table 1: Reductive cases for products of classical groups with at least one factor of type B or C.

$b(m),\ m \geq 2$  $H = \mathrm{SO}(2m),$ inside $\mathrm{SO}(2m+1)$
$b'(m),\ m \geq 2$  $H = \mathrm{N}\,\mathrm{SO}(2m),$ inside $\mathrm{SO}(2m+1)$
$b^{**}(3)$  $H = \mathsf{G}_2,$ inside $\mathrm{SO}(7)$
$c(m),\ m \geq 3$  $H = \mathrm{SL}(2) \times \mathrm{Sp}(2m-2),$ inside $\mathrm{Sp}(2m)$
$aa(1,1) + c^*(m),\ m \geq 2$  $H = \mathrm{SL}(2) \times \mathrm{Sp}(2m-2),$ inside $\mathrm{SL}(2) \times \mathrm{Sp}(2m)$ via the map $(X,Y) \mapsto (X, X \oplus Y)$
$b^{**}(3) + b^*(4)$  $H = \mathrm{N}\,\mathrm{SO}(7),$ inside $\mathrm{SO}(9)$
$a(1) + c^*(m),\ m \geq 2$  $H = \mathbb{C}^* \times \mathrm{Sp}(2m-2),$ inside $\mathrm{Sp}(2m)$
$a'(1) + c^*(m),\ m \geq 2$  $H = \mathrm{N}(\mathbb{C}^* \times \mathrm{Sp}(2m-2)),$ inside $\mathrm{Sp}(2m)$
$bx(3,2) + c^*(m),\ m \geq 2$  $H = \mathrm{Sp}(2m) \times \mathrm{Sp}(4),$ inside $\mathrm{Sp}(2m+4) \times \mathrm{Sp}(4)$ via the map $(X,Y) \mapsto (X \oplus Y, Y)$
$bx(3,2)$ the localization of $bx(3,2) + c^*(m)$ in $\{\alpha_1, \alpha_2, \alpha_3; \alpha'_1, \alpha'_2\}$ $H = \mathrm{Sp}(4)$ diagonally embedded in $\mathrm{SL}(4) \times \mathrm{Sp}(4)$
$bx(2,1,1) + c^*(m),\ m \geq 2$ the localization of $bx(3,2) + c^*(m)$ in $\{\alpha_1; \alpha_3, \alpha_4, \ldots, \alpha_{m+3}; \alpha'_1, \alpha'_2\}$ $H = \mathrm{Sp}(2m-2) \times \mathrm{SL}(2) \times \mathrm{SL}(2),$ inside $\mathrm{Sp}(2m) \times \mathrm{SL}(2) \times \mathrm{Sp}(4)$ via the map $(X,Y,Z) \mapsto (X \oplus Y, Z, Y \oplus Z)$
$bx(2,1,1)$ the localization of $bx(3,2) + c^*(m)$ in $\{\alpha_1; \alpha_3; \alpha'_1, \alpha'_2\}$ $H = \mathrm{SL}(2) \times \mathrm{SL}(2),$ inside $\mathrm{Sp}(4) \times \mathrm{SL}(2) \times \mathrm{SL}(2)$ via the map $(X,Y) \mapsto (X \oplus Y, X, Y)$

Table 1: Reductive cases for products of classical groups with at least one factor of type B or C (continuation).

$aa(1,1)+c^*(m)+c^*(n),\ m,n\geq 2$  $H=\mathrm{Sp}(2m-2)\times\mathrm{SL}(2)\times\mathrm{Sp}(2n-2)$, inside $\mathrm{Sp}(2m)\times\mathrm{Sp}(2n)$ via the map $(X,Y,Z)\mapsto(X\oplus Y,Y\oplus Z)$
$ax(1,1,1)+c^*(l)+c^*(m)+c^*(n),\ l,m,n\geq 2$  $H=\mathrm{Sp}(2l-2)\times\mathrm{Sp}(2m-2)\times\mathrm{Sp}(2n-2)\times\mathrm{SL}(2)$, inside $\mathrm{Sp}(2l)\times\mathrm{Sp}(2m)\times\mathrm{Sp}(2n)$ via the map $(X,Y,Z,W)\mapsto(X\oplus W,Y\oplus W,Z\oplus W)$
$ax(1,1,1)+c^*(l)+c^*(m),\ l,m\geq 2$ the localization of $ax(1,1,1)+c^*(l)+c^*(m)+c^*(n)$ in $\{\alpha_1,\ldots,\alpha_l;\alpha'_1,\ldots,\alpha_m;\alpha''_1\}$ $H=\mathrm{Sp}(2l-2)\times\mathrm{Sp}(2m-2)\times\mathrm{SL}(2)$, inside $\mathrm{Sp}(2l)\times\mathrm{Sp}(2m)\times\mathrm{SL}(2)$ via the map $(X,Y,W)\mapsto(X\oplus W,Y\oplus W,W)$
$ax(1,1,1)+c^*(l),\ l\geq 2$ the localization of $ax(1,1,1)+c^*(l)+c^*(m)+c^*(n)$ in $\{\alpha_1,\ldots,\alpha_l;\alpha'_1;\alpha''_1\}$ $H=\mathrm{Sp}(2l-2)\times\mathrm{SL}(2)$, inside $\mathrm{Sp}(2l)\times\mathrm{SL}(2)\times\mathrm{SL}(2)$ via the map $(X,W)\mapsto(X\oplus W,W,W)$
$cx(3,2)$  $H=\mathrm{Sp}(4)\times\mathrm{SL}(2)$, inside $\mathrm{Sp}(4)\times\mathrm{Sp}(6)$ via the map $(X,Y)\mapsto(X,X\oplus Y)$
$bx(2,1,1)+c^*(m)+c^*(n),\ m,n\geq 2$  $H=\mathrm{SL}(2)\times\mathrm{SL}(2)\times\mathrm{Sp}(2m-2)\times\mathrm{Sp}(2n-2)$, inside $\mathrm{Sp}(4)\times\mathrm{Sp}(2m)\times\mathrm{Sp}(2n)$ via the map $(X,Y,Z,W)\mapsto(X\oplus Y,X\oplus Z,Y\oplus W)$
$cc(p,p),\ p\geq 3$  $H=$ the normalizer of $\mathrm{Sp}(2p)$ diagonally embedded in $\mathrm{Sp}(2p)\times\mathrm{Sp}(2p)$
$bb(p,p),\ p\geq 2$  $H=\mathrm{SO}(2p+1)$ diagonally embedded in $\mathrm{SO}(2p+1)\times\mathrm{SO}(2p+1)$
$cc(p+q),\ p\geq 2$ even, $q\geq 2$  $H=\mathrm{Sp}(p+2q-2)\times\mathrm{Sp}(p+2),$ inside $\mathrm{Sp}(2p+2q)$

Table 1: Reductive cases for products of classical groups with at least one factor of type B or C (continuation).

$cc'(p+2),\ p\geq 2$ even  $H=\mathrm{N}(\mathrm{Sp}(p+2)\times\mathrm{Sp}(p+2)),$ inside $\mathrm{Sp}(2p+4)$
$ca'(m),\ m\geq 3$  $H=\mathrm{N}\,\mathrm{GL}(m),$ inside $\mathrm{Sp}(2m)$
$ca(m),\ m\geq 3$  $H=\mathrm{GL}(m),$ inside $\mathrm{Sp}(2m)$
$ba'(m),\ m\geq 3$  $H=\mathrm{N}\,\mathrm{GL}(m),$ inside $\mathrm{SO}(2m+1)$
$ba(m),\ m\geq 3$  $H=\mathrm{GL}(m),$ inside $\mathrm{SO}(2m+1)$
$a(1)+b'(m),\ m\geq 1$  $H=\mathbb{C}^*\times\mathrm{SO}(2m+1),$ inside $\mathrm{SO}(2m+3)$
$bo(p+q),\ p,q\geq 1$  $H=\mathrm{N}(\mathrm{SO}(p+1)\times\mathrm{SO}(p+2q)),$ inside $\mathrm{SO}(2(p+q)+1)$
$ay(2,1)+c^*(m),\ m\geq 2$  $H=\mathrm{SL}(2)\times\mathbb{C}^*\times\mathrm{Sp}(2m-2),$ inside $\mathrm{SL}(3)\times\mathrm{Sp}(2m)$ via the map $(X,a,Y)\mapsto(a^2\oplus a^{-1}X,X\oplus Y)$
$ax(1+p+1,1)+c^*(m),\ p\geq 1,\ m\geq 2$  $H=\mathrm{SL}(p+1)\times\mathbb{C}^*\times\mathrm{SL}(2)\times\mathrm{Sp}(2m-2),$ inside $\mathrm{SL}(p+3)\times\mathrm{Sp}(2m)$ via the map $(X_1,a,X_2,Y)\mapsto(a^{p+1}X_1\oplus a^{-2}X_2,X_2\oplus Y)$
$bx(3,3)$  $H=\mathrm{SO}(6)$ diagonally embedded in $\mathrm{SO}(6)\times\mathrm{SO}(7)$

Table 1: Reductive cases for products of classical groups with at least one factor of type B or C (continuation).

$dx(p,p),\ p \geq 4$  $H = \mathrm{SO}(2p)$ diagonally embedded in $\mathrm{SO}(2p+1) \times \mathrm{SO}(2p)$
$dx(p-1,p),\ p \geq 4$ the localization of $dx(p,p)$ in $\{\alpha_2, \ldots, \alpha_p; \alpha'_1, \ldots, \alpha'_p\}$ $H = \mathrm{SO}(2p-1)$ diagonally embedded in $\mathrm{SO}(2p-1) \times \mathrm{SO}(2p)$

Let us now prove existence for the systems in Table 1 which are neither tail cases, see §3.4, nor *strict*, see [3]. It is enough to analyze the following few cases: we can skip systems of rank < 3, which are well known, and systems that are localizations of others.

First of all, we examine the cases with defect 0.

- $bx(3,2) + c^*(m)$: Despite the presence of the spherical root $c^*(m)$, this case is not a tail one: there is no subset of colors Δ', in the notation of §3.4.

  The candidate subgroup H is equal to its normalizer, hence wonderful: call $\mathscr{S}$ the corresponding spherical system, with set of spherical roots Σ. The group H is contained in the subgroup $\mathrm{Sp}(2m) \times \mathrm{Sp}(4) \times \mathrm{Sp}(4)$ (inside $\mathrm{Sp}(2m+4) \times \mathrm{Sp}(4)$ by means of $(X, Y, Z) \mapsto (X \oplus Y, Z)$) and this induces a morphism to its wonderful embedding, which has the rank 2 system $ac(3) + c(m)$ (times a trivial factor under the action of $\mathrm{Sp}(4)$):

  

  From this morphism we deduce that $\alpha_1, \ldots, \alpha_{m+2} \in \mathrm{supp}\Sigma$.

  Moreover, $\mathscr{S}$ admits no factorization and so we have that α'_1 and/or α'_2 is in $\mathrm{supp}\Sigma$. The two colors moved by α_2 and by α_4 on the system $ac(3) + c(m)$ must lift to two colors on $\mathscr{S}$. If we examine all primitive systems having defect 0 with the help of these considerations, we easily see that $\mathscr{S}$ must be $bx(3,2) + c^*(m)$.
- $cx(3,2)$: Since $H = \mathrm{N}_G H$, then H is wonderful. It is included in $H' = \mathrm{Sp}(4) \times \mathrm{SL}(2) \times \mathrm{Sp}(4)$ (by means of $(X,Y) \mapsto (X, Y, X)$) and H' in $G \cong \mathrm{Sp}(4) \times \mathrm{Sp}(6)$ (by means of $(X, Y, Z) \mapsto (X, Y \oplus Z)$). The spherical system of the wonderful embedding of G/H' has diagram

  

  and this assures that $\alpha'_1, \alpha'_2, \alpha'_3 \in \mathrm{supp}\Sigma$.

  Since H is indecomposable, there must be some spherical root having support also on α_1 and/or α_2. Also, H is not a parabolic induction so one concludes that both α_1 and α_2 are in $\mathrm{supp}\Sigma$. We exclude the presence of any $a_1 \times a_1$ spherical root and the presence of any $c^*(m)$ spherical root because the only chances for them to appear would lead to the system $aa(1,1) + c^*(m) + c^*(n)$ (already known) or to a system where the defect is non-zero. So $\Sigma = S$. If we examine all primitive systems having defect 0 with the help of these considerations, we easily see that the corresponding spherical system must be $cx(2,3)$.

- $dx(p,p)$, $p \geq 4$: Again, $H = \mathrm{N}_G H$ so H is wonderful. It is diagonally included in $H' = \mathrm{SO}(2p) \times \mathrm{SO}(2p)$ and H' in $G = \mathrm{SO}(2p+1) \times \mathrm{SO}(2p)$ (by means of $\mathrm{SO}(2p) \subset \mathrm{SO}(2p+1)$). The spherical system of the wonderful embedding of G/H' has the following diagram.



Therefore, this case is similar, one can deduce that $\Sigma = S$ and again the condition $\mathrm{d} = 0$ forces the system to be $dx(p,p)$.

The cases $dx(p-1,p)$, $bx(3,3)$, $bx(3,2)$, $bx(1,1,2)$ are localizations of $dx(p,p)$.

Let us now discuss the two remaining cases: they both have defect 1.

- $ca(m)$: The candidate subgroup is spherically closed. One can see G/H as the set of couples of m-dimensional subspaces of $\mathbb{C}^{2m}$ satisfying the conditions:

$$G/H = \big\{(\pi_1, \pi_2) \mid \pi_1, \pi_2 \subset \mathbb{C}^{2m},\ \dim \pi_i = m,\ \pi_1 \cap \pi_2 = 0,\ J|_{\pi_i} = 0\big\}$$

where J is the bilinear form defining the group $\mathrm{Sp}(2m)$. If B is the Borel subgroup stabilizing the flag $\ldots \subset V_i \subset V_{i+1} \subset \ldots$ of subspaces of $\mathbb{C}^{2m}$ with $\dim V_i = i$, then the conditions:

$$\pi_1 \cap V_m \neq 0, \quad \pi_2 \cap V_m \neq 0$$

give two colours which are swapped by the action of the group $\mathrm{N}_G H/H \cong \mathbb{Z}/2\mathbb{Z}$. Moreover, these two colours are moved only by α_m so $\alpha_m \in \Sigma$. If one localizes on $S' = \{\alpha_1, \ldots, \alpha_{m-1}\}$ then one obtains a localization with defect equal to zero, and this forces the system of H to be $ca(m)$.
- $ba(m)$: This case is similar to $ca(m)$.

4.2. **Non-tail cases in type C.** For G containing components of type C, there are non-reductive primitive cases which are not tail cases, but do contain a spherical root of type $c^*(m)$: there is no subset of colors Δ' with the required properties, in the notations of §3.4. Also, for them L doesn't contain the typical factor $\mathrm{Sp}(2m-2)$ found in tail cases. We give as an example the following case:

- $ae_7(5) + c^*(m)$:



The subsets of colors Δ_{Q_-} and Δ_K defined in §3.1 are both equal to $\{D_1^+, D_1^-, D_2^+, D_2^-, D_4^-, D_6\}$, therefore $Q_- = K = P_{-S\setminus\{\alpha_3\}}$.



It follows that H has Levi component $L = L_K = L_{Q_-}$ isogenous to $\mathrm{GL}(3) \times \mathrm{Sp}(2m+2)$, and the unipotent radical of H is the unique L-submodule of K^u where $\mathrm{Sp}(2m+2)$ acts trivially.

The existence proof goes as follows. The subgroup H is wonderful, since $H = \mathrm{N}_G H$; it is included in K, and also in the wonderful subgroup $\mathrm{Sp}(6) \times \mathrm{Sp}(2m+2)$. The latter is cuspidal of has rank one, with spherical root $c^*(m+4)$. The wonderful variety X with generic stabilizer H is hence cuspidal, and there are four colors each moved by exactly one of the simple roots $\alpha_2, \alpha_3, \alpha_4, \alpha_6$. This forces S^p_X by dimension reasons, and hence Σ_X is determined. The family $\mathbf{A}_X$ follows easily from these considerations and $\mathrm{d}(X) = 0$.

4.3. **Primitive cases in type** $\mathsf{B} - \mathsf{C}$**.** In Table 2 we list some primitive spherical systems, without quotients of higher defect and without tails, for G a product of classical groups with at least one factor of type B or C. The remaining spherical systems (primitive, without quotients of higher defect and without tails, in type $\mathsf{B} - \mathsf{C}$) can all be obtained by localization from those listed in the table.

For each system, we describe the generic stabilizer H, together with a parabolic subgroup Q_- with compatible Levi decompositions: $H = L\,H^u$ and $Q_- = L_{Q_-}\,Q^u_-$ with $H^u \subset Q^u_-$ and $L = L_{Q_-} \cap H$. Here the connected center of L is always equal to that of L_{Q_-}, due to the absence of quotients with higher defect.

We denote by $V_L(\gamma)$ the simple L-submodule of $\mathrm{Lie}\,Q^u_-$ of highest weight γ. To simplify notations, we write highest weights for L-modules as T-weights, which must be intended as implicitly restricted to $T \cap L$. When it is more convenient, we describe the L-module Q^u_-/H^u instead of H^u.

Table 2: Primitive cases for products of classical groups with at least one factor of type B or C.

$bc^*(m),\ m \geq 2$ even  $Q_- = P_{-\{\alpha_1,\dots,\alpha_{m-1}\}}$ $(L,L) = \mathrm{Sp}(m)$ $H^u = (Q^u_-, Q^u_-)$
$ac^*(m) + c^*(n),\ m \geq 2$ even, $n \geq 2$  $Q_- = P_{-\{\alpha_2,\dots,\alpha_{m+n-1}\}}$ $(L,L) = \mathrm{Sp}(m+2n-4) \times \mathrm{Sp}(m)$ inside L_{Q_-} via the map $(X,Y) \mapsto (X \oplus Y)$ $Q^u_-/H^u = \mathbb{C}^m_Y$
$aa(1+1+1) + c^*(m),\ m \geq 2$  $Q_- = P_{-\{\alpha_1,\alpha_3,\dots,\alpha_{m+2}\}}$ $L = L_{Q_-}$ $H^u = (Q^u_-, Q^u_-)$

Table 2: Primitive cases for products of classical groups with at least one factor of type B or C. (continuation)

$by(4)$  $Q_- = P_{-\{\alpha_1,\alpha_3,\alpha_4\}}$ $(L,L) = \mathrm{SL}(2)\times\mathrm{SL}(2)$ inside L_{Q_-} via the map $(X,Y)\mapsto(X,X\oplus Y)$ $Q_-^u/H^u = \mathbb{C}^2_X$
$ce_7(5)$  $Q_- = P_{-\{\alpha_1,\alpha_2,\alpha_4,\alpha_5\}}$ $L = L_{Q_-}$ $H^u = (Q_-^u, Q_-^u)$
$bw(3)$  $Q_- = P_{-\{\alpha_1,\alpha_3\}}$ $(L,L) = \mathrm{SL}(2)$ inside L_{Q_-} via the map $X\mapsto(X,X)$ $Q_-^u/H^u = \mathbb{C}^2_X$
$\text{2-comb} + a(p),\ p\geq 1$  $Q_- = P_{-\{\alpha_1,\ldots,\alpha_p\}}$ $L = L_{Q_-}$ $H^u = V_L(-\alpha_{p+2})\oplus V_L(-\alpha_1-2\alpha_2-\ldots-2\alpha_{p+2})$
$ax(1+p+1+1+1)+c^*(m),\ p\geq 1, m\geq 2$  $Q_- = P_{-\{\alpha_1,\ldots,\alpha_p,\alpha_{p+2},\alpha_{p+4},\ldots,\alpha_{p+m+3}\}}$ $L = L_{Q_-}$ $H^u = (Q_-^u, Q_-^u)$
$ax(1+p+1,1)+a(q),\ p,q\geq 1$  $Q_- = P_{-\{\alpha_1,\ldots,\alpha_p,\alpha_{p+2},\ldots,\alpha_{p+q+1},\alpha_{p+q+3}\}}$ $L = L_{Q_-}$ $H^u = V_L(-\alpha_1-\ldots-\alpha_{p+q+1})\oplus$ $V_L(-\alpha_1-\ldots-\alpha_{p+1}-2\alpha_{p+2}-\ldots-2\alpha_{p+q+2}-\alpha_{p+q+3})\oplus$ $V_L(-2\alpha_1-\ldots-2\alpha_{p+q+2}-\alpha_{p+q+3})\oplus$ $V_L(-2\alpha_{p+2}-\ldots-2\alpha_{p+q+2}-\alpha_{p+q+3})$

Table 2: Primitive cases for products of classical groups with at least one factor of type B or C. (continuation)

$ax(1+p+1,1)+a(1),\ p\geq 1$  $Q_-=P_{-\{\alpha_1,\alpha_3,\ldots,\alpha_{p+2},\alpha_{p+4}\}}$ $L=L_{Q_-}$ $H^u=V_L(-\alpha_3-\ldots-\alpha_{p+4})\oplus V_L(-\alpha_1-\ldots-\alpha_{p+1}-2\alpha_{p+2}-2\alpha_{p+3}-\alpha_{p+4})\oplus$ $V_L(-2\alpha_1-\ldots-2\alpha_{p+3}-\alpha_{p+4})\oplus V_L(-2\alpha_{p+2}-2\alpha_{p+3}-\alpha_{p+4})$
$bx(3,2)+a(q),\ q\geq 1$  $Q_-=P_{-\{\alpha_1,\ldots,\alpha_{q+2},\alpha_{q+4},\alpha_{q+5}\}}$ $L=L_{Q_-}$ $H^u=(Q^u_-,Q^u_-)$
$ay(2+1+1)+c^*(m),\ m\geq 2$  $Q_-=P_{-\{\alpha_2,\alpha_4,\ldots,\alpha_{m+3}\}}$ $L=L_{Q_-}$ $H^u=(Q^u_-,Q^u_-)$
$ay(2,1)+a(q),\ q\geq 1$  $Q_-=P_{-\{\alpha_2,\ldots,\alpha_{q+1},\alpha_{q+3}\}}$ $L=L_{Q_-}$ $H^u=H^u=V_L(-\alpha_1-\ldots-\alpha_{q+1})\oplus$ $V_L(-\alpha_1-2\alpha_2-\ldots-2\alpha_{q+2}-\alpha_{q+3})\oplus$ $V_L(-2\alpha_1-\ldots-2\alpha_{q+2}-\alpha_{q+3})\oplus$ $V_L(-2\alpha_2-\ldots-2\alpha_{q+2}-\alpha_{q+3})$
$ay(2,1)+a(1)$  $Q_-=P_{-\{\alpha_2,\alpha_4\}}$ $L=L_{Q_-}$ $H^u=V_L(-\alpha_1-\alpha_2-2\alpha_3-\alpha_4)\oplus V_L(-\alpha_3-\alpha_4)\oplus$ $V_L(-2\alpha_1-2\alpha_2-2\alpha_3-\alpha_4)\oplus V_L(-2\alpha_3-\alpha_4)$
$ay(2,1)+b^*(2)$  $Q_-=P_{-\{\alpha_1,\alpha_3,\alpha_4\}}$ $(L,L)=\mathrm{SL}(2)\times\mathrm{SL}(2)$ inside L_{Q_-} via the map $(X,Y)\mapsto(X,X\oplus Y)$ $Q^u_-/H^u=\mathbb{C}^2_Y$

Table 2: Primitive cases for products of classical groups with at least one factor of type B or C. (continuation)

$by(p,p)+a(q),\ p\geq 2, q\geq 1$  $Q_-=P_{-\{\alpha_1,\ldots,\alpha_{2p+q-1}\}}$ $(L,L)=\mathrm{SL}(p+q)\times\mathrm{SL}(p)$ inside L_{Q_-} via the map $(X,Y)\mapsto(X\oplus Y)$ $Q_-^u/H^u=\mathbb{C}^{p+q}_X$
$by(p,p-1)+a(q),\ p\geq 2, q\geq 1$  $Q_-=P_{-\{\alpha_1,\ldots,\alpha_{2p+q-2}\}}$ $(L,L)=\mathrm{SL}(p+q-1)\times\mathrm{SL}(p)$ inside L_{Q_-} via the map $(X,Y)\mapsto(X\oplus Y)$ $Q_-^u/H^u=\mathbb{C}^{p}_Y$
$ae_7(5)+c^*(m),\ m\geq 2$  $Q_-=P_{-\{\alpha_1,\alpha_2,\alpha_4,\ldots,\alpha_{m+4}\}}$ $L=L_{Q_-}$ $H^u=(Q_-^u,Q_-^u)$

Let us discuss a few examples from Table 2.

- $by(p,p)+a(q)$. We have $Q_-=P_{-\{\alpha_1,\ldots,\alpha_{2p+q-1}\}}$, and the quotient by Δ_K is the following parabolic induction of $aa((p-1)+(q+1)+(p-1))$:

  

  It follows that (L,L) is equal to $\mathrm{SL}(p+q)\times\mathrm{SL}(p)$ inside $(L_{Q_-},L_{Q_-})=\mathrm{SL}(2p+q)$ via the map $(X,Y)\mapsto X\oplus Y$. The unipotent radical H^u has codimension $p+q$ inside Q_-^u, and this determines H^u uniquely, in such a way that $Q_-^u/H^u=\mathbb{C}^{p+q}_X$. To prove that this gives a geometric realization of $by(p,p)+a(q)$, we observe that H is wonderful and its spherical system $\mathscr{S}$ is not a parabolic induction. It has only two quotients, induced by the inclusions $H\subset K$ and $H\subset Q_-$, thus $\mathscr{S}$ is primitive. Then $\alpha_1,\ldots,\alpha_p,\alpha_{p+q+1},\ldots,\alpha_{2p+q}$ must be spherical roots, with α_i and $\alpha_{2p+q+1-i}$ having a color in common, for $i=1,\ldots,p$. The other colors in **A** follow immediately; dimensions and the inclusion $H\subset K$ force $\mathscr{S}$ to be $by(p,p)+a(q)$.

- $ay(2,1)+b^*(2)$ and $by(4)$. These two cases have same defect, same dimension, and they both admit a quotient equal to:

  

  which is parabolic induction of the rank 2 reductive case for $\mathrm{SL}(2)\times\mathrm{Sp}(4)$ with generic stabilizer $\mathrm{SL}(2)\times\mathrm{SL}(2)$, embedded via the map $(X,Y)\mapsto(X,X\oplus Y)$. This is also the commutator of the Levi part L of H. The parabolic subgroup Q_- is $P_{-\{\alpha_1,\alpha_3,\alpha_4\}}$, and it remains to find H^u inside Q_-^u, a L-submodule of codimension 2.

  The $(\mathrm{SL}(2)\times\mathrm{SL}(2))$-module Q_-^u is $\mathbb{C}\oplus\mathbb{C}^2_X\oplus\mathbb{C}^2_Y\oplus(\mathrm{Sym}^2\mathbb{C}^2_X\otimes\mathbb{C}^2_Y)$, where (Q_-^u,Q_-^u) is the summand $\mathbb{C}$. Therefore H^u may be obtained cutting $\mathbb{C}^2_X$ or $\mathbb{C}^2_Y$. The two choices will give two different generic stabilizers, both wonderful. It is then easy to show that they must correspond to our two systems. To decide which is which, it is enough to observe that $ay(2,1)+b^*(2)$ admits also a quotient to the rank 1 case:

  

  which gives an inclusion $H\subset SO(8)$.

4.4. **Minimal cases of projective fibrations.** For any minimal case of §3.2 we report a generic stabilizer H with notations as in §4.3. We denote by $V_M(\gamma)$ the simple M-submodule of $\mathrm{Lie}\,Q_-^u$ of highest weight γ, where M can be L or L_{Q_-}.

The first two cases are of rank two and already known.

(1) H is reductive, equal to the Levi factor of the parabolic subgroup $Q_-=P_{-\{\alpha_2,\dots,\alpha_n\}}$. By the way, this is a tail case already discussed in 3.4.
(2) Q_- is equal to $P_{-\{\alpha_2\}}$, L is equal to L_{Q_-} and $\mathrm{Lie}\,H^u$ is equal to $V_L(-3\alpha_1-\alpha_2)$.
(3) Q_- is equal to $P_{-\{\alpha_1,\alpha_2,\alpha_3\}}$, L is $\mathbb{C}^*\times\mathrm{SL}(4)$ and $\mathrm{Lie}\,H^u$ is the complementary of $V_L(-\alpha_4)\oplus V_L(-\alpha_1-2\alpha_2-3\alpha_3-2\alpha_4)$.
(4) Q_- is equal to $P_{-\{\alpha_1,\alpha_2\}}$, L is $\mathbb{C}^*\times\mathrm{GL}(3)$ and $\mathrm{Lie}\,H^u$ is the complementary of $V_L(-\alpha_3)\oplus V_L(-\alpha_3-\alpha_4)\oplus V_L(-\alpha_4)$.
(5) Q_- is equal to $P_{-\{\alpha_2\}}$, L is $(\mathbb{C}^*)^2\times\mathrm{SL}(2)$ and $\mathrm{Lie}\,H^u$ is the direct sum of simple L-modules as follows: $\mathfrak{m}$ diagonal in $V_L(-\alpha_1)\oplus V_L(-\alpha_3)$, plus $(\mathfrak{m},\mathfrak{m})$, plus all the simple modules in $\mathrm{Lie}\,P^u$ except $V_L(-\alpha_1)$, $V_L(-\alpha_3)$, $V_L(-\alpha_4)$, $V_L(-\alpha_1-\alpha_2-\alpha_3-\alpha_4)$ and $V_L(-\alpha_2-2\alpha_3-\alpha_4)$.
(6)
   (a) If n is even, Q_- is $P_{-\{\alpha_1,\dots,\alpha_{n-1}\}}$, L is $\mathbb{C}^*\times\mathrm{Sp}(n)$ and $\mathrm{Lie}\,H^u$ is just the simple L-submodule of codimension 1 in $V_{L_{Q_-}}(-\alpha_{n-1}-2\alpha_n)$.
   (b) If n is odd, Q_- is $P_{-\{\alpha_2,\dots,\alpha_{n-1}\}}$, L is $\mathbb{C}^*\times\mathrm{Sp}(n-1)$ and $\mathrm{Lie}\,H^u$ is the direct sum of four simple L-modules $\oplus_{i=1}^4\mathfrak{m}_i$. The module $\mathfrak{m}_1$ is diagonal in $V_{L_{Q_-}}(-\alpha_1)\oplus V_{L_{Q_-}}(-\alpha_n)$. The module $\mathfrak{m}_2$ is equal to $(\mathfrak{m}_1,\mathfrak{m}_1)$. The module $\mathfrak{m}_3$ is a codimension 1 L-submodule of $V_{L_{Q_-}}(-\alpha_{n-1}-2\alpha_n)$. The module $\mathfrak{m}_4$ is equal to $V_{L_{Q_-}}(-\alpha_1-2\alpha_2-\ldots-2\alpha_n)$.

REFERENCES

[1] P. Bravi, *Wonderful varieties of type E*, Represent. Theory **11** (2007) 174–191.
[2] P. Bravi, *Primitive spherical systems*, arXiv:0909.3765v1 .
[3] P. Bravi, S. Cupit-Foutou, *Classification of strict wonderful varieties*, arXiv:0806.2263v1 .

[4] P. Bravi, D. Luna, *An introduction to wonderful varieties with many examples of type F4*, arXiv:0812.2340v2 .

[5] P. Bravi, G. Pezzini, *Wonderful varieties of type D*, Represent. Theory **9** (2005) 578–637.

[6] M. Brion, *Classification des espaces homogènes sphériques*, Compos. Math. **63** (1987), 189–208.

[7] S. Cupit-Foutou, *Invariant Hilbert schemes and wonderful varieties*, arXiv:0811.1567v2 .

[8] S. Cupit-Foutou, *Wonderful varieties: a geometrical realization*, arXiv:0907.2852v1 .

[9] I.V. Losev, *Proof of the Knop conjecture*, Ann. Inst. Fourier (Grenoble) **59** (2009) 1105–1134.

[10] I.V. Losev, *Uniqueness property for spherical homogeneous spaces*, Duke Math. J. **147** (2009) 315–343.

[11] D. Luna, *Toute variété magnifique est sphérique*, Transform. Groups **1** (1996) 249–258.

[12] D. Luna, *Variété sphériques de type A*, Publ. Math. Inst. Hautes Études Sci. **94** (2001) 161–226.

[13] G. Pezzini, *Wonderful varieties of type C*, Ph.D. Thesis, Università La Sapienza, Rome, 2004.